



ELSEVIER

Mechanisms of Development 52 (1995) 399–400



Volume contents

Volume 52 (1995)

Issue contents

D.H. Rowitch, A.P. McMahon

Pax-2 expression in the murine neural plate precedes and encompasses the expression domains of *Wnt-1* and *En-1*

M. Hollyday, J.A. McMahon, A.P. McMahon

Wnt expression patterns in chick embryo nervous system

A.F. Candia, C.V.E. Wright

The expression pattern of *Xenopus Mox-2* implies a role in initial mesodermal differentiation

A. Nichols, E. Rungger-Brändle, L. Muster, D. Rungger

Inhibition of *Xhox1A* gene expression in *Xenopus* embryos by antisense RNA produced from an expression vector read by RNA polymerase III

B. Leussink, A. Brouwer, M. El Khattabi, R.E. Poelmann, A.C. Gittenberger-de Groot, F. Meijlink

Expression patterns of the paired-related homeobox genes *MHox/Prx1* and *S8/Prx2* suggest roles in development of the heart and the forebrain

M. Tirronen, V.-P. Lahti, T.I. Heino, C. Roos

Two *otu* transcripts are selectively localised in *Drosophila* oogenesis by a mechanism that requires a function of the *otu* protein

S.S. Sathe, P.J. Harte

The *Drosophila extra sex combs* protein contains WD motifs essential for its function as a repressor of homeotic genes

T.R. Breen, V. Chinwalla, P.J. Harte

Trithorax is required to maintain *engrailed* expression in a subset of *engrailed*-expressing cells

M.A. van Dijk, L.T.C. Peltenburg, C. Murre

Hox gene products modulate the DNA binding activity of Pbx1 and Pbx2

1 **K. Verschuereen, N. Dewulf, M.-J. Goumans, O. Lonnay, A. Feijen, S. Grimsby, K. Vande Spiegle, P. ten Dijke, A. Morén, P. Vanscheeuwijck, C.-H. Heldin, K. Miyazono, C. Mummery, J. Van Den Eijnden-Van Raaij, D. Huylebroeck**

3 Expression of type I and type IB receptors for activin in midgestation mouse embryos suggests distinct functions in organogenesis 109

9 U. Ernsberger, H. Patzke, J.P. Tissier-Seta, T. Reh, C. Goridis, H. Rohrer

The expression of tyrosine hydroxylase and the transcription factors cPhox-2 and Cash-1: evidence for distinct inductive steps in the differentiation of chick sympathetic precursor cells 125

C.K. Motzny, R. Holmgren

The *Drosophila* cubitus interruptus protein and its role in the *wingless* and *hedgehog* signal transduction pathways 137

Issue contents 151

A.R. Ungar, G.M. Kelly, R.T. Moon

Wnt4 affects morphogenesis when misexpressed in the zebrafish embryo 153

J.C. Richardson, D. Gatherer, H.R. Woodland

Developmental effects of over-expression of normal and mutated forms of a *Xenopus* NF- κ B homologue 165

K.J. Blake, B. Rogina, A. Centurion, S.L. Helfand

Changes in gene expression during post-eclosion development in the olfactory system of *Drosophila melanogaster* 179

A.K. Cheng, E.J. Robertson

The murine LIM-kinase gene (*limk*) encodes a novel serine threonine kinase expressed predominantly in trophoblast giant cells and the developing nervous system 187

V. Fridmacher, M. Le Bert, F. Guillo, S. Magre

Switch in the expression of the K19/K18 keratin genes as a very early evidence of testicular differentiation in the rat 199

M.J. Stassen, D. Bailey, S. Nelson, V. Chinwalla, P.J. Harte

The *Drosophila trithorax* proteins contain a novel variant of the nuclear receptor type DNA binding domain and an ancient conserved motif found in other chromosomal proteins 209

S.S. Sathe, P.J. Harte

The *extra sex combs* protein is highly conserved between *Drosophila virilis* and *Drosophila melanogaster* 225

A. Fjose, U. Weber, M. Mlodzik

A novel vertebrate *svp*-related nuclear receptor is expressed as a step gradient in developing rhombomeres and is affected by retinoic acid 233

C.M. Turner, P.N. Adler

Morphogenesis of *Drosophila* pupal wings in vitro 247

D. Ferrari, L. Sumoy, J. Gannon, H. Sun, A.M.C.

Brown, W.B. Upholt, R.A. Kosher

The expression pattern of the *Distal-less* homeobox-containing gene *Dlx-5* in the developing chick limb bud suggests its involvement in apical ectodermal ridge activity, pattern formation, and cartilage differentiation 257

M. Reichman-Fried, B.-Z. Shilo

Breathless, a *Drosophila* FGF receptor homolog, is required for the onset of tracheal cell migration and tracheole formation 265

Y.-Q. Wang, A. Sizeland, X.-F. Wang, D. Sassoon

Restricted expression of type-II TGF β receptor in murine embryonic development suggests a central role in tissue modeling and CNS patterning 275

E.A. Argao, M.J. Kern, W.W. Branford, W.J. Scott, Jr., S.S. Potter

Malformations of the heart, kidney, palate, and skeleton in α -MHC-*Hoxb-7* transgenic mice 291

S.-G. Gong, B.A. Reddy, L.D. Etkin

Two forms of *Xenopus* nuclear factor 7 have overlapping spatial but different temporal patterns of expression during development 305

J. Ellis, Q. Liu, M. Breitman, N.A. Jenkins, D.J. Gilbert, N.G. Copeland, H.V. Tempest, S. Warren, E. Muir, H. Schilling, F.A. Fletcher, S.F. Ziegler, J.H. Rogers

Embryo brain kinase: a novel gene of the *eph/ek* receptor tyrosine kinase family 319

M.-O. Fauvarque, V. Zuber, J.-M. Dura

Regulation of *polyhomeotic* transcription may involve local changes in chromatin activity in *Drosophila* 343

J.H. Clement, P. Fettes, S. Knöchel, J. Lef, W. Knöchel

Bone morphogenetic protein 2 in the early development of *Xenopus laevis* 357

S. Rinkwitz-Brandt, M. Justus, I. Oldenettel, H.-H. Arnold, E. Bober

Distinct temporal expression of mouse *Nkx-5.1* and *Nkx-5.2* homeobox genes during brain and ear development 371

G. Conway

A novel gene expressed during zebrafish gastrulation identified by differential RNA display 383

Author index 393

Subject index 395

Volume contents 399

